

QUAIL AND ORNAMENTAL POULTRY FEED

Starter and maturing feed for quail and ornamental poultry

- deuka quail and ornamental poultry feeds (= W/Z) are suitable for rearing and maturing quails, pheasants, partridges, guinea fowl, peacocks and ornamental geese and ducks.
- The micro-granulated W/Z starter optimally covers the high protein and amino acid requirements in the rearing phase.
- deuka W/Z feed provides your quails and ornamental poultry with all the nutrients and minerals following the chick phase until they are ready to lay or, in the case of fattening birds, until they are slaughtered.
- Our W/Z feeds do **not** contain **any** coccidiostats.

Ingredients

W/Z Starter:

Crude protein	26,0 %
Crude fat	4,50 %
Crude fibre	3,20 %
Crude ash	5,40 %
Lysine	1,50 %
Methionine	0,50 %
Calcium	1,30 %
Phosphorus	1,00 %
Sodium	0,15 %
Energy	11.40 MJ ME/kg

W/Z feed:

Crude protein	18,0 %
Crude fat	4,50 %
Crude fibre	3,20 %
Crude ash	5,40 %
Lysine	0,85 %
Methionine	0,40 %
Calcium	1,00 %
Phosphorus	0,60 %



Sodium	0,15 %
Energy	12.20 MJ ME/kg

For individual values, there may be slight deviations in content between the supplier plants.

List of ingredients

W/Z Starter

Soybean extruded meal, steam fermented, wheat, maize, wheat semolina bran, monocalcium phosphate, vegetable fatty acids, calcium carbonate, sodium chloride

W/Z feed:

Wheat, soybean extruded meal, steam, maize, wheat semolina bran, calcium carbonate, vegetable fatty acids, monocalcium phosphate, sodium chloride

Depending on the production plant, season, availability of goods, etc., slight changes in composition may be possible. The actual recipe and ingredients of the feed can be found on the declaration attached to each feed bag.

Vitamins

Vitamin A	10.000.00 I.U./kg
Vitamin D3	3.000.00 I.U./kg
Vitamin E	50.00 mg/kg

For individual values, there may be slight deviations in content between the supplier plants.

Trace elements

contain other trace elements

For individual values, there may be slight deviations in content between the supplier plants.

Feeding recommendation

W/Z Starter:

Feed as starter feed for quails, pheasants, guinea fowl, partridges and peacocks in the first 5-6 weeks of life. Quails up to the 3rd week of life. If W/Z Starter is to be used for ornamental geese and ducks, it is recommended to use up to max. 2 weeks of life.

W/Z feed:

As a complete feed for quails, pheasants, partridges, guinea fowls, peacocks, ornamental ducks and geese. To be used following the respective rearing feed as maturity/maintenance feed or until laying maturity. To be used for fattening animals until slaughter.

Pheasant chicks as well as other ornamental poultry and quails feed on notched animals in the wild and therefore make very high demands on the chick feed. They are grain eaters and prefer granulated or pelleted feed. Cleanliness and feed hygiene are top priorities for these sensitive animals. The partial intolerance to some coccidiostats must be taken into account. Therefore, the **deuka W/Z feeds** do not contain any coccidiostats.

Always provide fresh drinking water.

What, when, how much - the feeding recommendation

	Feed type	Quantity
Breeding 1st - 6th week (quails up to 3rd week)	deuka W/Z Starter	for free intake
Maturity / Growth / Fattening following the respective rearing feed	deuka W/Z Mature feed	for free intake

deuka

Laying phase quails	deuka laying quail feed	approx. 30 g / bird and day
Laying phase other ornamental poultry	deuka all-mash L (cooked) or all-mash breeding (cooked) or deuka laying meal (cooked) deuka grains extra in a ratio of 2:1	depending on the type of poultry and weight between approx. 120 g (pheasants) and approx. 300 g (peacocks) per bird and day